

- Prieto-Fingerhut T, Banovac K, Lynne CM. A study comparing sterile and nonsterile urethral catheterization in patients with spinal cord injury. *Rehabil Nurs* 1997; 22: 299–302
- Reynard JM, Vass J, Sullivan ME et al. Sphincterotomy and the treatment of detrusor-sphincter dyssynergia: current status, future prospects. *Spinal Cord* 2003; 41: 1–11
- Riedl CR, Stephen RL, Daha LK et al. Electromotive administration of intravesical bethanechol and the clinical impact on acontractile detrusor management: introduction of a new test. *J Urol* 2000; 164: 2108–2111
- Roe B, Williams K, Palmer M. Bladder training for urinary incontinence in adults. *Cochrane Database Syst Rev* 2000; 2: CD001308
- Schurch B, de Seze M, Denys P et al. Botulinum toxin type a is a safe and effective treatment for neurogenic urinary incontinence: results of a single treatment, randomized, placebo controlled 6-month study. *J Urol* 2005; 174: 196–200
- Schurch B, Rodic B, Jeanmonod D. Posterior sacral rhizotomy and intradural anterior sacral root stimulation for treatment of the spastic bladder in spinal cord injured patients. *J Urol* 1997; 157: 610–614
- Schurch B, Schmid DM, Stohrer M. Treatment of neurogenic incontinence with botulinum toxin A. *Neur Urol J* 2000; 342: 665
- Stöhrer M, Castro-Diaz D, Chartier-Kastler E et al. EAU guidelines on neurogenic lower urinary tract dysfunction. http://www.uroweb.org/fileadmin/user_upload/Guidelines/neurogenic.pdf. 2003: 1–40
- Stone AR. Neurourologic evaluation and urologic management of spinal dysraphism. *Neurosurg Clin N Am* 1995; 6: 269–277
- Swierzewski SJ 3rd, Gormley EA, Belville WD et al. The effect of terazosin on bladder function in the spinal cord injured patient. *J Urol* 1994; 151: 951–954
- van Kerrebroeck P, Abrams P, Lange R et al. Duloxetine versus placebo in the treatment of European and Canadian women with stress urinary incontinence. *BJOG* 2004; 111: 244–251
- van Kerrebroeck P, Rezapour M, Cortesse A et al. Desmopressin in the treatment of nocturia: a double-blind, placebo-controlled study. *Eur Urol* 2007; 52: 221–229
- van Kerrebroeck PE, Koldewijn EL, Rosier PF et al. Results of the treatment of neurogenic bladder dysfunction in spinal cord injury by sacral posterior root rhizotomy and anterior sacral root stimulation. *J Urol* 1996; 155: 1378–1381
- Yamanishi T, Yasuda K, Homma Y et al. A multicenter placebo-controlled, double-blind trial of urapidil, an alpha-blocker, on neurogenic bladder dysfunction. *Eur Urol* 1999; 35: 44–51
- Yasuda K, Yamanishi T, Kawabe K et al. The effect of urapidil on neurogenic bladder: a placebo controlled double-blind study. *J Urol* 1996; 156: 1125–1130

Aus: Hans-Christoph Diener, Christian Weimar (Hrsg.)

Leitlinien für Diagnostik und Therapie in der Neurologie

Herausgegeben von der Kommission "Leitlinien" der Deutschen Gesellschaft für Neurologie

Thieme Verlag, Stuttgart, September 2012

© Deutsche Gesellschaft für Neurologie