

- 23: 706–716
- Rietberg MB, Brooks D, Uitdehaag BM et al. Exercise therapy for multiple sclerosis. *Cochrane Database Syst Rev* 2005; 1: 1–26
 - Saunders DH, Greig CA, Mead GE et al. Physical fitness training for stroke patients. *Cochrane Database Syst Rev* 2009; 4: CD003316
 - Scheidtmann K, Fries W, Müller F et al. Effect of levodopa in combination with physiotherapy on functional motor recovery after stroke: a prospective, randomised, double-blind study. *Lancet* 2001; 358: 787–790
 - Sirtori V, Corbetta D, Moja L et al. Constraint-induced movement therapy for upper extremities in stroke patients. *Cochrane Database Syst Rev* 2009; 4: CD004433
 - Sterr A. Training-based interventions in motor rehabilitation after stroke: theoretical and clinical considerations. *Behav Neurol* 2004; 15: 55–63
 - Steultjens EM, Dekker J, Bouter LM et al. Occupational therapy for stroke patients: a systematic review. *Stroke* 2003; 34: 676–687
 - Stinear C. Prediction of recovery of motor function after stroke. *Lancet Neurol* 2010; 9: 1228–1232
 - van Peppen RP, Kwakkel G, Wood-Dauphinee S et al. The impact of physical therapy on functional outcomes after stroke: what's the evidence? *Clin Rehabil* 2004; 18: 833–862
 - van Swigchem R, Vloothuis J, den Boer J et al. Is transcutaneous peroneal stimulation beneficial to patients with chronic stroke using an ankle-foot orthosis? A within-subjects study of patients' satisfaction, walking speed and physical activity level. *J Rehabil Med* 2010; 42: 117–121
 - van Vliet PM, Lincoln NB, Foxall A. Comparison of Bobath based and movement science based treatment for stroke: a randomised controlled trial. *J Neurol Neurosurg Psychiatry* 2005; 76: 500–503
 - Ward NS, Frackowiak RSJ. The functional anatomy of cerebral reorganisation after focal brain damage. *J Physiol (Paris)* 2006; 99: 425–236
 - WHO (World Health Organization). Internationale Klassifikation der Funktionsfähigkeit, Behinderung und Gesundheit (ICF). Stand Oktober 2005. Herausgegeben vom Deutschen Institut für Medizinische Dokumentation und Information, DIMDI. WHO-Kooperationszentrum für das System Internationaler Klassifikationen (www.dimdi.de; 11.01.11)
 - Wolf SL, Winstein CJ, Miller JP et al. Effect of constraint-induced movement therapy on upper extremity function 3 to 9 months after stroke. The EXCITE Randomized Clinical Trial. *J Am Med Assoc* 2006; 296: 2095–2104
 - Wu HM, Tang JL, Lin XP et al. Acupuncture for stroke rehabilitation. *Cochrane Database Syst Rev* 2006; 3: CD004131

Aus: Hans-Christoph Diener, Christian Weimar (Hrsg.)
Leitlinien für Diagnostik und Therapie in der Neurologie
Herausgegeben von der Kommission "Leitlinien" der Deutschen Gesellschaft für
Neurologie
Thieme Verlag, Stuttgart, September 2012

© Deutsche Gesellschaft für Neurologie